

Editorial Board

Yoshiro Abe, Tokushima University, Japan

Noriko Aramaki, Keio University, Japan

Jun Arata, National Hospital Organization Kyoto Medical Center, Japan

Masaki Fujioka, National Hospital Organization Nagasaki Medical Center, Japan

Toshihiro Fujiwara, Hyogo College of Medicine, Japan

Hiroshi Furukawa, Aichi Medical University, Japan

Koichi Gonda, Tohoku Medical and Pharmaceutical University, Japan

Kazunobu Hashikawa, Kobe University, Japan

Ichiro Hashimoto, Tokushima University, Japan

Akiteru Hayashi, Toho University, Japan

Takayuki Honda, Iwate Medical University, Japan

Hideki Kadota, Kyushu University, Japan

Hisakazu Kato, Gifu University, Japan

Yoshihiro Kimata, Okayama University, Japan

Tateki Kubo, Osaka University, Japan

Daisuke Maeda, Osaka University, Japan

Hajime Matsumura, Tokyo Medical University, Japan

Takeo Matsusue, Kansai Electric Power Hospital, Japan

Kyoichi Matsuzaki, International University of Health and Welfare, Japan

Hiroshi Mizuno, Juntendo University, Japan

Hisashi Motomura, Osaka City University, Japan

Takashi Nakatsuka, JR Tokyo General Hospital, Japan

Toshiaki Numajiri, Kyoto Prefectural University, Japan

Rei Ogawa, Nippon Medical School, Japan

Norihiko Ohura, Kyorin University, Japan

Takafumi Ohshiro, Keikoh-kai Ohshiro Clinic, Japan

Takaharu Otsuka, Showa University, Japan

Nobuhiro Sato, Showa University, Japan

Ichiro Tanaka, Tokyo Dental College, Japan

Katsumi Tanaka, Nagasaki University, Japan

Rica Tanaka, Juntendo University, Japan

Koichi Tomita, Osaka University, Japan

Kazuhiro Toriyama, Nagoya City University, Japan

Yasuyoshi Tosa, Showa University, Japan

Tetsuji Uemura, Saga University, Japan

Takashi Yamaki, Tokyo Women's Medical University, Japan

Kotaro Yoshimura, Jichi Medical University, Japan